

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5






Fig. 6






Title: PRETARGETING METHODS AND COMPOUNDS

					<u></u>					
	+PBS	SD	+NON-BT	SD	+10:1	SD	+5:1	SD	BT-SAT'D	SD
BLOOD	31.05		29.94	1.35	8.54	0.91	7.03	0.18	24.58	0.68
TAIL	2.43		1.80	0.09	1.46	0.09	1.76	0.04	1.96	0.40
FUNG	1.47		1.09	0.22	0.54	0.10	0.48	0.07	0.76	0.01
LIVER	5.42	0.69	4.66	0.36	9.60	1.20	9.11	0.41	97.9	90.0
SPLEEN	0.25		0.34	0.03	0.17	0.03	0.18	0.00	0.38	0.02
STOMACH	0.28		0.33	0.03	0.53	0.34	0.49	0.00	0.29	0.04
KIDNEY	1.72		1.38	0.08	2.76	0.00	3.28	0.32	1.58	0.08
INTESTINE	3.40		3.44	0.10	4.22	0.02	6.62	0.14	2.83	0.13
	46.02		42.98		27.83		28.95		39.13	
	Group 1		Group 2		Group 3		Group 4		Group 5	
										•

Fig. 8



% IN1 DOZE IN Brood







Fig. 11B










Fig. 15A




Fig. 15B




Fig. 16A




Fig. 16B




Fig. 17A




Fig. 17B




Fig. 18




Fig. 19




Fig. 20